



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61M	A2	(11) International Publication Number: WO 99/51284 (43) International Publication Date: 14 October 1999 (14.10.99)
<p>(21) International Application Number: PCT/US99/07061</p> <p>(22) International Filing Date: 31 March 1999 (31.03.99)</p> <p>(30) Priority Data: 60/080,658 3 April 1998 (03.04.98) US</p> <p>(71) Applicant (for all designated States except US): MALLINCK-RODT INC. [US/US]; 675 McDonnell Boulevard, P.O. Box 5840, St. Louis, MO 63134 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): RAJAGOPALAN, Raghavan [US/US]; 13031 Vinson Court, Maryland Heights, MO 63043 (US). BUGAJ, Joseph, E. [US/US]; 2916 Kettering Drive, St. Charles, MO 63303 (US). DORSHOW, Richard, B. [US/US]; 11977 Niehaus Lane, St. Louis, MO 63146 (US). ACHILEFU, Samuel, I. [NG/US]; 3424 San Sevilla Court, St. Louis, MO 63044 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published Without international search report and to be republished upon receipt of that report.</p>
<p>(54) Title: NON-COVALENT BIOCONJUGATES USEFUL FOR DIAGNOSIS AND THERAPY</p>		
<p>(57) Abstract</p> <p>A composition comprising non-covalent carrier-hapten bioconjugates having the formula HM—CM wherein HM is a hapten molecule whose molecular weight is generally, but not always, less than 1000 Daltons and is capable of performing specific functions; CM is a carrier molecule, whose molecular weight is generally, but not always, more than 1000 Daltons and is capable of transporting the hapten to a specific site; and the dashed line is a non-covalent bond between the carrier molecule and the hapten molecule. Preferably, the bioconjugates are formed from fluorescent dye haptens such as cyanine, indocyanine, squaraine, porphyrins, Rose Bengal, and methylene blue dye and carrier molecules such as methylated serum albumin, polyarginine, polyaspartic acid, polyglutamic acid, cyclodextrin, and inulin. The bioconjugates are useful in diagnostic and therapeutic medical procedures because they are stable <i>in vitro</i> before being used and stable <i>in vivo</i> during and after use.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

NON-COVALENT BIOCONJUGATES USEFUL FOR DIAGNOSIS AND THERAPY

BACKGROUND OF THE INVENTION

Field of the Invention

5 This invention pertains to non-covalent-carrier-hapten bioconjugates and their use to selectively deliver diagnostic and therapeutic agents to target tissues and organs. Particularly, this invention pertains to the selective delivery of fluorescent non-covalent carrier-hapten bioconjugates to a specific site for the purpose of determining the structure and function of tissues or organs using optical tomographic
10 imaging procedures, fluorescence monitoring procedures, absorbance monitoring procedures, or endoscopic examination procedures.

Description of the Prior Art

Non-covalent intermolecular forces (e.g. electrostatic, hydrogen bonding and Van der Waals interactions) play a vital role in many biological process such as
15 enzyme catalysis, drug-receptor interaction, antigen-antibody interaction, biotin-avidin interaction, DNA double helix formation, phagocytosis, pigmentation in plant and animals, and cellular transport. Variety of colors observed in flowers and plants, for instance, are attributed to non-covalent association between the natural pigments and carbohydrates or proteins found in plant cells.

20 Non-covalent forces can alter the physicochemical and/or biological properties of haptens or carriers. For example, association of dye or a pigment molecule with proteins or carbohydrates can change the chemical or photo stability, change the intensity and/or the wavelength of absorption/emission maxima, or both. Although the interaction energy per unit interaction is quite small (c.a. less than 40
25 kJ/interaction), the cumulative effect of multiple points of interaction along the two surfaces can be substantial and can lead to strong binding between the hapten and the carrier. This approach has been successfully used to prepare anti-DNA antibodies. DNA is a highly charged anionic macromolecule that is normally non-immunogenic; but, when it is complexed with a highly charged cationic methylated bovine serum

albumin (MBSA), DNA becomes immunogenic. Thus, the non-covalent DNA-MBSA bioconjugate was stable enough to elicit immune response toward DNA.

Non-specific interactions also play an important role in biological processes. For example, human serum albumin binds various molecules in a non selective fashion and facilitates the transport of these molecules across from the vasculature to the cells.

Thus, it is clear from the examples above that the non-covalently attached bioconjugates function biologically as a single unit. In addition to the above mentioned properties, the carrier molecule may also protect some haptens from chemical, photochemical, or radiolytic degradation. The present invention is intended to exploit the concept of non-covalent interactions in the design of novel bioconjugates for diagnosis and therapy.

Further, the use of fluorescent dyes and covalent dye bioconjugates for the detection of various species in the body fluids is well known in the field of in-vitro immunodiagnosics. Dye-immunoconjugates have been widely used in immunohistology, and immunochemical detection of various antigens, hormones, drugs and the like in body fluids via radioimmunoassay (RIA) or enzyme-linked immunosorbent assay (ELISA) procedures. Fluorescent dyes have also been used as chemosensors and in fluorescence microscopy. (J.P. Desvergne and A.W. Czarnik, *Chemosensors of Ion and Molecule Recognition*, Kluwer Academic, Boston, 1997; B.H. Satir, Ed., *Noninvasive Techniques in Cell Biology*, Wiley-Liss, New York, 1990). Indocyanine green, a highly fluorescent polyene dye, has been used for monitoring cardiac output, assessing hepatic function, and tomographic imaging of tumors (K. Licha *et al.*, Synthesis and Characterization of Cyanine Dyes as Contrast Agents for Near-Infrared Imaging, *SPIE*, 1996, Vol. 2927, 192-197; X. Li *et al.*, Tumor Localization Using Fluorescence of Indocyanine Green in Rat Models, *SPIE*, 1995, Vol. 2389, 789-798; B. Riefke *et al.*, In Vivo Characterization of Cyanine Dyes as Contrast Agents for Near-Infrared Imaging, *SPIE*, 1996, Vol. 2927, 199-208).

In conventional bioconjugate chemistry, conjugates are prepared by covalent attachment of various effector molecules such as drugs, hormones,

radiopharmaceutical agents, magnetic resonance imaging agents, chemotherapeutic agents, and the like to the bioactive carriers. Such a process often involves cumbersome chemical manipulation of the two components, in addition to the complicated synthesis of appropriate activated haptens necessary for covalent attachment. Moreover, the bioactivity of the resulting complex is, in many cases, either greatly diminished or obviated altogether.

Polyene dyes, in general, and indocyanine green, in particular, have several disadvantages that limit their use as near-infrared (NIR) contrast agents. Indocyanine green has a very short plasma half-life and is rapidly taken up by the liver (D.K Meijer *et al.*, Pharmacokinetics of Biliary Excretion in Man. VI. Indocyanine Green, *Eur. J. Clin. Pharmacol.*, 1988, Vol. 35, 295-303); has low fluorescence efficiency, and undergoes degradation in aqueous media with loss of fluorescence. Therefore, there is a need for novel bioconjugates that are simple to prepare and stable enough to be useful as a diagnostic or a therapeutic agent.

15

SUMMARY OF THE INVENTION

It is, therefore, an object of the present invention to provide non-covalent carrier-hapten bioconjugates that are simple to prepare and are useful in diagnostic and therapeutic medical procedures.

It is another object of the present invention to provide bioconjugates that are stable during storage before being used in diagnostic and therapeutic medical procedures.

It is a further object of the present invention to provide a method for performing a diagnostic or therapeutic procedure on a patient.

It is a another object of the present invention to provide a method for altering the blood persistence of a hapten.

It is a another object of the present invention to provide a method for increasing the *in vitro* and *in vivo* fluorescent life of a fluorescent dye.

These and other objects are achieved using new and structurally diverse non-covalent carrier-hapten bioconjugates having the formula:



wherein HM is a hapten molecule whose molecular weight is generally, but not
5 always, less than 1000 Daltons and is capable of performing specific functions; CM is
a carrier molecule, whose molecular weight is generally, but not always, more than
1000 Daltons and is capable of transporting the hapten to a specific site; and the
dashed line is a non-covalent bond between the carrier molecule and the hapten
molecule.

10 Preferably, the bioconjugates are formed from fluorescent dye haptens and carrier
molecules selected from the group consisting of serum albumin, methylated serum
albumin, polypeptides with molecular weight ranges from 2000 to 20000 Daltons, and
polysaccharides with molecular weight ranges from 2000 to 20000 Daltons,
polynucleotides with molecular weight ranges from 2000 to 100000 Daltons,
15 cyclodextrins, calixarenes, and surfactants.

Most preferably, the bioconjugates are formed from fluorescent dye haptens
selected from the group consisting of cyanine, indocyanine, squaraine, porphyrins,
Rose Bengal, and methylene blue dye and carrier molecules selected from the group
consisting of methylated serum albumin, polyarginine, polyaspartic acid, polyglutamic
20 acid, cyclodextrin, inulin, polyadenylic acid, and polyguanylic acid.

The bioconjugates are useful as diagnostic and therapeutic agents in medical
procedures because they are stable *in vitro* before being used and stable *in vivo* during
and after use. It is theorized that the bond between the hapten and the carrier
stabilizes the bioconjugates and delays its degradation *in vitro* by environmental
25 factors. Also, the bond causes the bioconjugates to be metabolized slowly *in vivo* and
therefore retained in the body for a longer period than expected for the non-bound
hapten.

Other objects, advantages, and novel features of the present invention will
become apparent in the following detailed description of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 shows the rat blood clearance rate for indocyanine green dye.

Fig. 2 shows the rat blood clearance rate for an inulin-indocyanine green dye bioconjugate.

5 Fig. 3 shows the rat blood clearance rate for a MBSA-indocyanine green dye bioconjugate.

Fig. 4 shows the rat blood clearance rate for a gamma cyclodextrin-indocyanine green dye bioconjugate.

10 Fig. 5 shows the rat blood clearance rate for a polyaspartic acid-indocyanine green dye bioconjugate.

Fig. 6(a) shows the rat blood clearance rate for the polyaspartic-indocyanine green dye bioconjugate and for indocyanine green dye alone at the time of initial preparation in aqueous solution.


15 Fig. 6(b) shows the rat blood clearance rate for the polyaspartic-indocyanine green dye bioconjugate and for indocyanine green dye alone at one (1) day after preparation.

Fig. 6(c) shows the rat blood clearance rate for the polyaspartic-indocyanine green dye bioconjugate and for indocyanine green dye alone at five (5) days after preparation.

20 Fig. 6(d) shows the rat blood clearance rate for the polyaspartic-indocyanine green dye bioconjugate and for indocyanine green dye alone at nine (9) days after preparation.

DETAILED DESCRIPTION OF THE INVENTION

25 The present invention provides new and structurally diverse non-covalent carrier-hapten bioconjugates having the formula:



wherein HM is a hapten molecule whose molecular weight is generally, but not always, less than 1000 Daltons and is capable of performing specific functions; CM is a carrier molecule, whose molecular weight is generally, but not always, more than 1000 Daltons and is capable of transporting the hapten to a specific site; and the
5 dashed line is a non-covalent bond between the carrier molecule and the hapten molecule.

The hapten is either a small molecule or a macromolecule and is selected from the group consisting peptides, carbohydrates, and fluorescent dyes. The hapten can be molecules such as hormones, antibodies, anti-neoplastic agents, enzymes, coenzymes,
10 peptidomimetics, glycomimetics, cell adhesion molecules, radionuclide metal complexes, magnetic resonance imaging agents, X-ray opacification agents, and echogenic agents.

Preferably, the hapten is a dye whose absorption and emission maxima fall in 200-1200 nanometers (nm) range or a photosensitizer capable of transferring energy
15 to the tissues or to the other components inside or outside the cell. Most preferably, the hapten is a fluorescent dye with absorption and emission maxima in the region of 450-950 nm.

The carrier is selected from the group consisting of proteins such as albumins and globulins, glycoproteins, polypeptides, polysaccharides, polynucleotides, lipoproteins,
20 surfactants, and other natural or synthetic polymeric substances.

Preferably, the carrier is a macromolecule selected from the group consisting of proteins such as albumin or methylated albumin, glycoproteins such as antibodies or selectins, polysaccharides such as inulin or lectins, polynucleotides such as DNA or RNA, inclusion compounds such as cyclodextrins or calixarenes, and surfactants such
25 as Tween. Most preferably the carrier is a polysaccharide, polynucleotide, protein, polypeptide, or cyclodextrin.

In a preferred embodiment, a bioconjugate consists of a fluorescent dye hapten with absorption and emission maxima in the region of 450-950 nm and a

polysaccharide carrier held together by a non-covalent bond between the dye and the polysaccharide.

In another preferred embodiment, a dye bioconjugate consists of a fluorescent dye hapten with absorption and emission maxima in the region of 450-950 nm and a
5 protein held together by a non-covalent bond between the dye and the protein.

In a further preferred embodiment, a bioconjugate consists of a fluorescent dye hapten with absorption and emission maxima in the region of 450-950 nm and a polypeptide carrier held together by a non-covalent bond between the dye and the polypeptide.

10 The dye molecules of the present invention may vary widely depending on the contemplated application. For diagnostic imaging of areas of lesion, fluorescent dyes absorbing and emitting in the near infrared (NIR) region, i.e. 700-900 nm, are desirable. These may include, but not limited to, cyanines, indocyanines, or squaraines. For monitoring blood clearance or for endoscopic examination of lesions,
15 dyes absorbing and emitting in the region of 450-950 nm, preferably 600-900 nm, are desirable.

Similarly, the carrier molecules may also vary widely. For blood persistent agents, albumin or methylated serum albumin is preferable. For renal function measurements, a polysaccharide or anionic polypeptides are desirable. For
20 endoscopic examination of lesions, antibodies and antibody fragments are preferred.

The non-covalent carrier-hapten bioconjugates of this present invention can either be advantageously prepared by simply mixing the two components in an optimal stoichiometric proportions and administering an effective amount of this mixture contained in a pharmaceutically acceptable formulation into individuals either
25 systemically or locally to the organ or tissue to be studied. Alternatively, the bioconjugates can be isolated and stored by methods well known in the art. The novel bioconjugates of the present invention have broad clinical utility, which includes, but not limited to, diagnostic imaging of tumors, inflammation (both sterile and bacterial), and impaired vasculature; laser guided endoscopic examination of sites

of lesion; photodynamic and chemotherapy of tumors or infection, and dynamic organ function assessment.

The novel bioconjugates of this invention can be formulated into diagnostic or therapeutic compositions for enteral, parenteral, or oral administration. These compositions contain an effective amount of bioconjugates along with conventional pharmaceutical carriers and excipients appropriate for the type of administration contemplated. These compositions may also include stabilizing agents such as ascorbic or gentisic acid. For example, parenteral compositions advantageously contain a sterile aqueous solution or suspension of bioconjugates whose concentration ranges from about 1 μM to about 10 mM. Preferred parenteral formulations have a concentration of bioconjugates of 100 μM to 2 mM. Such solutions also may contain pharmaceutically acceptable buffers and, optionally, electrolytes such as sodium chloride. Formulations for enteral administration may vary widely as is well-known in the art. In general, such formulations are liquids which include an effective amount of bioconjugates in aqueous solution or suspension. Such enteral composition may optionally include buffers, surfactants, thixotropic agents, and the like. Compositions for oral administration may also contain flavoring agents and other ingredients for enhancing their organoleptic qualities.

The diagnostic compositions are administered in doses effective to achieve the desired diagnostic or therapeutic objective. Such doses may vary widely depending upon the particular bioconjugates employed, the organs or tissues to be examined, the equipment employed in the clinical procedure, and the like.

The present invention also provides a method of performing a diagnostic or therapeutic procedure on a patient for the purpose of determining the structure and function of tissues or organs. The method comprises administering the non-covalent carrier-hapten bioconjugates of the present invention to a patient, allowing the bioconjugates to become localized in or around a tissue or organ, and performing a diagnostic or therapeutic procedure such as a optical tomographic imaging procedures, fluorescence monitoring procedures, absorbance monitoring procedures,

and endoscopic examination procedures, preferably a fluorescence monitoring procedure on the patient.

5 The present invention also provides a method for altering the blood persistence of a hapten by forming the non-covalent carrier-hapten bioconjugates of the present invention. Some bioconjugates are more stable *in vivo* than the hapten alone and are cleared from the patient's blood at a slower rate than the hapten alone. The slow blood clearance rate provides more time for doing a diagnostic and therapeutic medical procedure because the hapten remains in the blood system of the patient for an extended period. Other bioconjugates are less stable *in vivo* than the hapten alone and are cleared from the patient's blood at a faster rate than the hapten alone. The faster blood clearance rate requires less time for doing a diagnostic and therapeutic medical procedure because the hapten remains in the blood system of the patient for a shorter period. This means that the patient can spend less time having the procedure and recuperating.

15 The present invention also provides a method for increasing the *in vitro* and *in vivo* fluorescent life of a fluorescent dye by forming the non-covalent carrier-hapten bioconjugates of the present invention. The bioconjugates retain the ability to fluoresce for extended periods when compared to the fluorescent dye alone. This longer fluorescent life permits the bioconjugates to be stored on the shelf for extended periods and then used for diagnostic and therapeutic medical procedures.


The present invention also provides fluorescent dye bioconjugates that have a longer *in vitro* and *in vivo* fluorescent life than the fluorescent dye alone. This longer fluorescent life permits the bioconjugates to be stored on the shelf for extended periods and then used for diagnostic and therapeutic medical procedures.

25 The following examples illustrate the specific embodiment of the invention described in this document. As would be apparent to skilled artisans, various changes and modifications are possible and are contemplated within the scope of the invention described.

EXPERIMENTAL PROCEDURES

Example 1

Blood clearance properties of indocyanine green dye



Indocyanine Green Dye

5 A 500 μ L portion of indocyanine green solution (0.8 mg/mL) was injected through the lateral tail vein of Sprague-Dawley or Fischer rats. Blood clearance of the dye bioconjugate was monitored by fluorescence signal emanating from the capillaries of the ear surface using a laser fluorescence apparatus (excitation, 780 nm; emission
10 830 nm). The time constant (τ) for indocyanine green blood clearance is 138 seconds (Fig. 1).

Example 2

Preparation and blood clearance properties of a inulin-indocyanine green dye bioconjugate

15 A solution of inulin (MW 5000, 30 mg) in water (5 mL) was treated with indocyanine green (4 mg) and the entire mixture was shaken vigorously for about 2 minutes. The blood clearance rate of this bioconjugate was determined using the same procedure described in Example 1. The time constant (τ) for blood clearance for this bioconjugate is 130 seconds (Fig. 2). The blood clearance rate for this bioconjugate is
20 approximately the same as that for indocyanine green.

Example 3**Preparation and blood clearance properties of a MBSA-indocyanine green dye bioconjugate**

A solution of methylated bovine serum albumin (MBSA) (MW 70000, 25 mg) in
5 water (5 mL) was treated with indocyanine green (4 mg) and the entire mixture was
shaken gently for about 5 minutes. The blood clearance rate of this bioconjugate was
determined using the same procedure described in Example 1. The time constant (τ)
for blood clearance for this bioconjugate is 224 seconds (Fig. 3). The blood clearance
rate of this bioconjugate is approximately two times slower than that of indocyanine
10 green.

Example 4**Preparation and blood clearance properties of a gamma cyclodextrin-indocyanine green dye bioconjugate**

A solution of γ -cyclodextrin (MW 1100, 11.2 mg) in water (5 mL) was treated
15 with indocyanine green (4 mg) and the entire mixture was shaken gently for 1 hour.
The blood clearance rate of this bioconjugate was determined using the same
procedure described in Example 1. The time constant (τ) for blood clearance for this
bioconjugate is 142 seconds (Fig. 4). The blood clearance rate of this bioconjugate is
approximately the same as that for indocyanine green.

Example 5**Preparation and blood clearance properties of a polyaspartic acid-indocyanine green dye bioconjugate**

A solution of polyaspartic acid (MW 6000, 82 mg) in water (10 mL) was treated
with indocyanine green (4 mg) and the entire mixture was shaken gently for about 5
25 minutes. The blood clearance rate of this bioconjugate was determined using the same
procedure described in Example 1. The time constant (τ) for blood clearance for this
bioconjugate is 91 seconds (Fig. 5). The blood clearance rate of this bioconjugate is
faster than that for indocyanine green.

Referring to Examples 1-5, the data shows that various carriers such as albumins,
30 globulins, glycoproteins, polypeptides, polysaccharides, polynucleotides, lipoproteins,
and surfactants interact with indocyanine green dye to form non-covalent carrier-

hapten bioconjugates and that the blood persistence of a hapten is altered when in the form of a bioconjugate.

Example 6

Stability and photophysical properties of several indocyanine green dye bioconjugates

An aqueous solution of indocyanine green (ICG) and the carriers, inulin, poly-d,l-aspartic acid, polyguanylic acid, MBSA, and γ -cyclodextrin were prepared and the fluorescence intensity was monitored at ambient temperature at various time intervals. Table 1 and Figs. 6(a), 6(b), 6(c), and 6(d) show the stability data of the bioconjugates.

Table 1
Percent decrease in fluorescence intensity at 822 nanometers

Indocyanine Green Dye	Percent Decrease in Fluorescence		
	2 Hours	4 Hours	24 Hours
Water	0	19	48
Inulin	45	45	45
Polyaspartic acid	0	0	0
γ -Cyclodextrin	0	100	-
MBSA	0	0	-
Polyaspartic acid/BSA	0	0	0
Polyguanylic acid	0	0	0

Referring to Figs. 6(a), 6(b), 6(c), and 6(d), the data show the stability enhancement achieved by the polyaspartic acid-indocyanine green dye bioconjugate over that of the indocyanine green dye alone. Referring to Table 1, the data show that some stability enhancement is also seen for bioconjugates involving MBSA and polyguanylic acid.

Example 7


Preparation of carboxyphloroglucinol-squaraine dye

A mixture of 2,4,6-trihydroxybenzoic acid (3.40 g, 0.02 mol) and squaric acid (1.14 g, 0.01 mol) in glacial acetic acid (15 mL) was heated under reflux for 4 hours.

The cherry-red solution was poured onto water (100 mL) and the purple dye was collected by filtration, washed well with copious water, and dried. The dye was suspended in water (100 mL) and heated to boiling for about 10 minutes, filtered hot, and dried. The dye was then used as such for further studies (A. Treibs and K. Jacob.,
 5 *Angew. Chem. Int. Ed. English*, 1965, 4, 694).

Example 8

Stability and photophysical properties of several carboxyphloroglucinol-squaraine dye bioconjugates



10

Carboxyphloroglucinol Squaraine Dye

An aqueous solution of carboxyphloroglucinol-squaraine dye and the carriers, inulin, poly-l-arginine, poly-d,l-aspartic acid, and γ -cyclodextrin were prepared and the fluorescence was temporally monitored at ambient temperature over 48 hours (λ_{ex} = 515, λ_{em} = 613). The results are shown in Table 2.

15

Table 2
Percent decrease in fluorescence intensity at 613 nanometers

Carboxyphloroglucinol Squaraine Dye	Percent Decrease in Fluorescence		
	2 Hours	4 Hours	24 Hours
Water	13	13	50
Inulin	0	0	5
Polyarginine	8	20	43
Polyaspartic acid	41	55	55
γ -Cyclodextrin	27	37	41

Referring to Table 2, the data show that various carriers such as polysaccharides, inclusion compounds, and polynucleotides interact with carboxyphloroglucinol squaraine dye non-covalent carrier-hapten bioconjugates and that the *in vitro* and *in vivo* fluorescent life of the fluorescent dye is increased when in the form of a

5 bioconjugate

Example 9

Preparation of benzothiazolim-squaraine dye


A mixture of 2-methylbenzothiazole (2.98 g, 0.020 mol) and bromoacetic acid (2.98 g, 0.022 mol) was heated to about 100 °C for 15 minutes. The dark reaction

10 mixture was poured onto acetone (100 mL) and the lavender colored solid was collected by filtration, washed with acetone, and dried. A mixture of this benzothiazolium salt (1.38 g, 5 mmol) and squaric acid (0.28 g, 2.5 mmol) in n-butanol/toluene (1:1, 15 mL) was heated under Dean-Stark conditions for 4 hours. The dark blue solution was poured onto water (100 mL) and the blue dye was

15 collected by filtration, washed well with copious acetone, and dried (S. Das *et al.*, *J. Phys. Chem.*, 1996, 100, 17310-17315).

Example 10

Stability and photophysical properties of benzothiazolim-squaraine dye bioconjugates



Benzothiazolium Squaraine Dye

An aqueous solution of benzothiazolim-squaraine dye and the carriers, inulin, polyarginine, poly-d,l-aspartic acid, polyguanydylic acid, and γ -cyclodextrin were prepared and the fluorescence was temporally monitored at ambient temperature over 48 hours. The results are shown in Table 3.

5


Table 3
Percent decrease in fluorescence intensity at 665 nanometers

Benzothiazolim Dye	Percent Decrease in Fluorescence		
	2 Hours	4 Hours	24 Hours
Water	92	64	46
Inulin	92	54	54
Polyarginine	0	0	47
Polyaspartic acid	0	0	0
γ -Cyclodextrin	33	41	41

Referring to Table 3, the data show that various carriers such as polysaccharides, inclusion compounds, and polynucleotides interact with benzothiazolium squaraine dye non-covalent carrier-hapten bioconjugates and that the *in vitro* and *in vivo* fluorescent life of the fluorescent dye is increased when in the form of a bioconjugate.

10

Example 11
Preparation of cyanine blue dye conjugates



Cyanine Blue Dye

A mixture of 2-methylbenzothiazole (2.98 g, 0.020 mol) and bromoacetic acid (2.98 g, 0.022 mol) was heated to about 100 °C for 15 minutes. The dark reaction mixture was poured onto acetone (100 mL) and the lavender colored solid was collected by filtration, washed with acetone, and dried. A mixture of this

15

benzothiazolium salt (1.44 g, 5 mmol) and malonaldehyde tetramethyl acetal (0.42 g, 2.5 mmol) in pyridine was heated under reflux for 4 hours. The dark blue solution was poured onto ether (500 mL) and the blue dye was collected by filtration, washed well with copious ether, and dried.

5

Example 12**Stability and photophysical properties of cyanine blue dye conjugates**

An aqueous solution of cyanine blue dye and the carriers, inulin, polyarginine, poly-d,l-aspartic acid, and γ -cyclodextrin were prepared and the fluorescence was temporally monitored at ambient temperature over 24 hours. The results are shown in

10 Table 4.

Table 4
Percent decrease in fluorescence intensity at 670 nanometers


Cyanine Blue Dye	Percent Decrease in Fluorescence		
	2 Hours	4 Hours	24 Hours
Water	20	42	99
Inulin	6	10	16
Polyarginine	0	0	0
Polyaspartic acid	73	73	78
γ -Cyclodextrin	0	9	9

Referring to Table 4, the data show that various carriers such as polysaccharides, inclusion compounds, and polynucleotides interact with cyanine blue dye non-covalent carrier-hapten bioconjugates and that the *in vitro* and *in vivo* fluorescent life of the fluorescent dye is increased when in the form of a bioconjugate.

Although the invention has been described with respect to specific modifications, the details thereof are not to be construed as limitations, for it will be apparent that various equivalents, changes, and modifications may resorted to without departing from the spirit and scope thereof, and it is understood that such equivalent embodiments are to be included therein.

What is claimed is:

1. A composition comprising non-covalent carrier-hapten bioconjugates having the formula:



5 wherein HM is a hapten molecule selected from the group consisting of peptides, carbohydrates, photosensitizers, and fluorescent dyes; CM is a carrier molecule selected from the group consisting of proteins, glycoproteins, polypeptides, polysaccharides, inclusion compounds, polynucleotides, lipoproteins, and surfactants; and the dashed line is a non-covalent bond between the carrier molecule and the
10 hapten molecule.

2. The composition of claim 1 wherein HM is selected from the group consisting of peptides containing 10 or less amino acids, oligosaccharides containing 10 or less furanose or pyranose moieties, photosensitizers with absorption and emission maxima in the range of 200-1200 nm, and fluorescent dyes with absorption and emission
15 maxima in the range of 200-1200 nm.

3. The composition of claim 1 wherein CM is selected from the group consisting of serum albumin, methylated bovine albumin, polypeptides having a molecular weight of from 2000 to 20000 Daltons, polysaccharides having a molecular weight of from 2000 to 20000 Daltons, polynucleotides having a molecular weight of from
20 2000 to 100000 Daltons, cyclodextrins, calixarenes, and surfactants.

4. The composition of claim 1 wherein HM is selected from the group consisting of fluorescent dyes with absorption and emission maxima between 450-950 nm.

5. The composition of claim 4 wherein CM is selected from the group consisting of serum albumin, methylated serum albumin, polypeptides with a molecular weight of
25 from 2000 to 20000 Daltons, and polysaccharides with a molecular weight of from 2000 to 20000 Daltons, polynucleotides, cyclodextrins, calixarenes, and surfactants.

6. The composition of claim 4 wherein HM is selected from the group consisting of cyanine, indocyanine, squaraine, porphyrins, Rose Bengal, and methylene blue dye.

7. The composition of claim 6 wherein CM is selected from the group consisting of methylated serum albumin, polyarginine, polyaspartic acid, polyglutamic acid, cyclodextrin, and inulin.

8. A method of performing a diagnostic or therapeutic procedure on a patient,
5 comprising:

administering to a patient a diagnostic or therapeutic effective amount of non-covalent carrier-hapten bioconjugates having the formula:



wherein HM is a hapten molecule selected from the group consisting of peptides,
10 carbohydrates, photosensitizers, and fluorescent dyes; CM is a carrier molecule selected from the group consisting of proteins, glycoproteins, polypeptides, polysaccharides, inclusion compounds, polynucleotides, lipoproteins, and surfactants; and the dashed line is a non-covalent bond between the carrier molecule and the hapten molecule;

15 allowing the bioconjugates to become localized in or around a tissue or organ; and

performing a diagnostic or therapeutic procedure on the patient.

9. The method of claim 8 wherein said procedure is selected from the group of optical tomographic imaging procedures, fluorescence monitoring procedures,
20 absorbance monitoring procedures, and endoscopic examination procedures.

10. The method of claim 8 wherein HM is selected from the group consisting of peptides containing 10 or less amino acids, oligosaccharides containing 10 or less furanose or pyranose moieties, photosensitizers with absorption and emission maxima in the range of 200-1200 nm, and fluorescent dyes with absorption and emission
25 maxima in the range of 200-1200 nm.

11. The method of claim 8 wherein CM is selected from the group consisting of serum albumin, methylated bovine albumin, polypeptides having a molecular weight of from 2000 to 20000 Daltons, polysaccharides having a molecular weight of from

2000 to 20000 Daltons, and polynucleotides having a molecular weight of from 2000 to 100000 Daltons, polynucleotides, cyclodextrins, calixarenes, and surfactants.

12. The method of claim 8 wherein HM is selected from the group consisting of fluorescent dyes with absorption and emission maxima between 450-950 nm.


5 13. The method of claim 8 wherein CM is selected from the group consisting of serum albumin, methylated serum albumin, polypeptides with a molecular weight of from 2000 to 20000 Daltons, and polysaccharides with a molecular weight of from 2000 to 20000 Daltons, polynucleotides, cyclodextrins, calixarenes, and surfactants.

10 14. The method of claim 8 wherein said procedure is a fluorescence monitoring procedure.

15 15. The method of claim 14 wherein HM is selected from the group consisting of cyanine, indocyanine, squaraine, porphyrins, Rose Bengal, and methylene blue dye.

16 16. The method of claim 15 wherein CM is selected from the group consisting of methylated serum albumin, polyarginine, polyaspartic acid, polyglutamic acid, cyclodextrin, and inulin.

17. A method of for altering the blood persistence of a hapten comprising forming non-covalent carrier-hapten bioconjugates having the formula:



20 wherein HM is a hapten molecule selected from the group consisting of peptides, carbohydrates, photosensitizers, and fluorescent dyes; CM is a carrier molecule selected from the group consisting of proteins, glycoproteins, polypeptides, polysaccharides, polynucleotides, lipoproteins, and surfactants; and the dashed line is a non-covalent bond between the carrier molecule and the hapten molecule.

25 18. The method of claim 17 wherein HM is selected from the group consisting of peptides containing 10 or less amino acids, oligosaccharides containing 10 or less furanose or pyranose moieties, photosensitizers with absorption and emission maxima in the range of 200-1200 nm, and fluorescent dyes with absorption and emission maxima in the range of 200-1200 nm.

30 19. The method of claim 17 wherein CM is selected from the group consisting of serum albumin, methylated bovine albumin, polypeptides having a molecular weight

of from 2000 to 20000 Daltons, polysaccharides having a molecular weight of from 2000 to 20000 Daltons, and polynucleotides having a molecular weight of from 2000 to 100000 Daltons, polynucleotides, cyclodextrins, calixarenes, and surfactants.


20. The method of claim 17 wherein HM is selected from the group consisting of
5 fluorescent dyes with absorption and emission maxima between 450-950 nm.

21. The method of claim 20 wherein CM is selected from the group consisting of serum albumin, methylated serum albumin, polypeptides with a molecular weight of from 2000 to 20000 Daltons, and polysaccharides with a molecular weight of from 2000 to 20000 Daltons, polynucleotides, cyclodextrins, calixarenes, and surfactants.

10 22. The method of claim 20 wherein HM is selected from the group consisting of cyanine, indocyanine, squaraine, porphyrins, Rose Bengal, and methylene blue dye.

23. The method of claim 22 wherein CM is selected from the group consisting of methylated serum albumin, polyarginine, polyaspartic acid, polyglutamic acid, cyclodextrin, and inulin.

15 24. A method of increasing the *in vitro* and *in vivo* fluorescent life of a fluorescent dye comprising forming non-covalent carrier-hapten bioconjugates having the formula:



wherein HM is a hapten molecule selected from the group consisting of fluorescent
20 dyes; CM is a carrier molecule selected from the group consisting of proteins, glycoproteins, polypeptides, polysaccharides, inclusion compounds, polynucleotides, lipoproteins, and surfactants; and the dashed line is a non-covalent bond between the carrier molecule and the hapten molecule.

25 25. The method of claim 24 wherein HM is selected from the group consisting of fluorescent dyes with absorption and emission maxima in the range of 200-1200 nm.

26. The method of claim 24 wherein CM is selected from the group consisting of serum albumin, methylated bovine albumin, polypeptides having a molecular weight of from 2000 to 20000 Daltons, polysaccharides having a molecular weight of from 2000 to 20000 Daltons, and polynucleotides having a molecular weight of from 2000
30 to 100000 Daltons, polynucleotides, cyclodextrins, calixarenes, and surfactants.

27. The method of claim 24 wherein HM is selected from the group consisting of fluorescent dyes with absorption and emission maxima between 450-950 nm.

28. The method of claim 27 wherein CM is selected from the group consisting of serum albumin, methylated serum albumin, polypeptides with a molecular weight of
5 from 2000 to 20000 Daltons, and polysaccharides with a molecular weight of from 2000 to 20000 Daltons, polynucleotides, cyclodextrins, calixarenes, and surfactants.

29. The method of claim 27 wherein HM is selected from the group consisting of cyanine, indocyanine, squaraine, porphyrins, Rose Bengal, and methylene blue dye.

30. The method of claim 29 wherein CM is selected from the group consisting of
10 methylated serum albumin, polyarginine, polyaspartic acid, polyglutamic acid, cyclodextrin, and inulin.

FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6a




FIG. 6b






FIG. 6C





(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 October 1999 (14.10.1999)

PCT

(10) International Publication Number
WO 99/51284 A3

(51) International Patent Classification⁶: **C07K 16/00**,
I/00, A61K 51/00, 49/00, G01N 33/566, 33/531, 33/533

[NG/US]: 3424 San Sevilla Court, St. Louis, MO 63044 (US).

(21) International Application Number: PCT/US99/07061

(22) International Filing Date: 31 March 1999 (31.03.1999)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/080,658 3 April 1998 (03.04.1998) US

(71) Applicant (for all designated States except US):
MALLINCKRODT INC. [US/US]: 675 McDonnell
Boulevard, P.O. Box 5840, St. Louis, MO 63134 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **RAJAGOPALAN, Raghavan** [US/US]: 13031 Vinson Court, Maryland Heights, MO 63043 (US). **BUGAJ, Joseph, E.** [US/US]: 2916 Kettering Drive, St. Charles, MO 63303 (US). **DORSHOW, Richard, B.** [US/US]: 11977 Niehaus Lane, St. Louis, MO 63146 (US). **ACHILEFU, Samuel, I.**

(81) Designated States (national): AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
17 January 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 99/51284 A3

(54) Title: NON-COVALENT BIOCONJUGATES USEFUL FOR DIAGNOSIS AND THERAPY

(57) Abstract: A composition comprising non-covalent carrier-hapten bioconjugates having the formula HM-----CM wherein HM is a hapten molecule whose molecular weight is generally, but not always, less than 1000 Daltons and is capable of performing specific functions; CM is a carrier molecule, whose molecular weight is generally, but not always, more than 1000 Daltons and is capable of transporting the hapten to a specific site; and the dashed line is a non-covalent bond between the carrier molecule and the hapten molecule. Preferably, the bioconjugates are formed from fluorescent dye haptens such as cyanine, indocyanine, squaraine, porphyrins, Rose Bengal, and methylene blue dye and carrier molecules such as methylated serum albumin, polyarginine, polyaspartic acid, polyglutamic acid, cyclodextrin, and inulin. The bioconjugates are useful in diagnostic and therapeutic medical procedures because they are stable *in vitro* before being used and stable *in vivo* during and after use.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/07061**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) : Please See Extra Sheet.

US CL : 530/391.3, 403, 807; 424/1.57, 9.34; 436/501, 543, 546

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530/391.3, 403, 807; 424/1.57, 9.34; 436/501, 543, 546

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X -- Y	US 4,806,488 A (BERGER, JR. et al) 21 February 1989, Abstract.	1 -- 1-30
X -- Y	BERGER et al. Immunochemistry of the Subunits of Transcarboxylase. J Biol Chem. Vol. 251, No. 22, pages 7021-7034, especially Materials and Methods.	1 -- 1-30
Y	LEE, S.H. Cytochemical Study of Estrogen Receptor in Human Mammary Cancer. Am J Clin Pathol. August 1978, Vol. 70, No. 2, pages 197-203, especially Materials and Methods.	1-30

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

22 MAY 1999

Date of mailing of the international search report

15 JUN 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

HANKYEL T. PARK, PH.D.

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/07061

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

C07K 16/00, 1/00; A61K 51/00, 49/00; G01N 33/566, 33/531, 33/533

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, DIALOG, MEDLINE, BIOSIS, EMBASE, CA SEARCH

search terms: carrier, hapten, bioconjugate, noncovalent, conjugate, indocyanine, squaraine, inulin, fluorescent, bovine, albumin, dye